

Weiterführende Literatur zu Brachflächen als Biodiversitätsinseln in der Agrarlandschaft

Wietzke, A. et al. (2020). Flower strips, conservation field margins and fallows promote the arable flora in intensively farmed landscapes: Results of a 4-year study. *Agriculture, Ecosystems & Environment* 304. <https://doi.org/10.1016/j.agee.2020.107142>

Meichtry-Stier, K. S., Duplain, J., Lanz, M., Lugrin, B. & Birrer, S. (2018). The importance of size, location, and vegetation composition of perennial fallows for farmland birds. *Ecology and Evolution* 8: 9270–9281. <https://onlinelibrary.wiley.com/doi/full/10.1002/ece3.4420>

Tscharntke, T., Batáry, P. & Dormann, C. F. (2011). Set-aside management: How do succession, sowing patterns and landscape context affect biodiversity? *Agriculture, Ecosystems & Environment* 143: 37–44. <https://www.sciencedirect.com/science/article/pii/S0167880911000259>

Winkler, K.; Wackers, F., Bukovinszkine-Kiss, G., van Lenteren, J. (2006). Sugar resources are vital for Diadegma semiclausum fecundity under field conditions, *Basic and Applied Ecology*, Volume 7, Issue 2: 133-140. <https://www.sciencedirect.com/science/article/pii/S1439179105000770>

BMUV Entwurf NBS 2030: <https://dialog.bmuv.de/bmu/de/home/file/ fileId/810/name/Ziele-%20&%20Ma%C3%9Fnah-menkatalog%20zur%20NBS%202030.pdf>