

## Schweinswal – *Phocoena phocoena*

### Zitierte Literatur

- BEINEKE, A., SIEBERT, U., MCLACHLAN, M., BRUHN, P., THRON, K., FAILING, K., MÜLLER, G. & BAUMGÄRTNER, W. (2005): Investigations of the potential influence of environmental contaminants on the thymus and spleen of harbour porpoises (*Phocoena phocoena*). – Environmental Science & Toxicology 39: 3933-3938.
- BERGGREN, P., ISHAQ, R., ZEBÜHR, Y., NÄF, C., BAND, C. & BROMAN, D. (1999): Patterns and levels of organochlorines (DDTs, PCBs, non-ortho PCBs and PCDD/Fs) in male harbour porpoises (*Phocoena phocoena*) from the Baltic Sea, the Kattegat-Skagerrak Seas and the west coast of Norway. – Marine Pollution Bulletin 12: 1070-1084.
- BERGGREN, P., WADE, P. R., CARLSTRÖM, J. & READ, A. J. (2002): Potential limits to anthropogenic mortality for harbour porpoises in the Baltic region. – Biological Conservation 103: 313-322.
- BJØRGE, A. & TOLLEY, K. A. (2009): Harbour porpoise *Phocoena phocoena*. – In: PERRIN, W. F., WÜRSIG, B. & THEWISSEN, J. G. M. (Hrsg.): Encyclopedia of Marine Mammals, Second Edition. – San Diego/ USA (Academic Press): 530-533.
- BRÄGER, S. & ZWEIFEL, U. L. (2009): Noise Pollution. – In: HELSINKI COMMISSION (Hrsg.): Biodiversity in the Baltic Sea. – Baltic Sea Environment Proceedings No. 116B: 125-128.
- BRANDT, M. J., DIEDERICHS, A., BETHKE, K. & NEHLS, G. (2011): Responses of harbour porpoises to pile driving at the Horns Rev II offshore wind farm in the Danish North Sea. – Marine Ecology Progress Series 421: 205-216.
- BRUHN, R., KANNAN, N., PETRICK, G., SCHULZ-BULL, D. E. & DUINKER J. C. (1999): Persistent chlorinated organic contaminants in harbour porpoises from the North Sea, the Baltic Sea and Arctic waters. – The Science of the Total Environment 237/238: 351-361.
- BUNDESAMT FÜR SEESCHIFFFAHRT UND HYDROGRAPHIE (BSH): Leitsätze für die Anwendung der Eingriffsregelung innerhalb der ausschließlichen Wirtschaftszone und auf dem Festlandsockel im Rahmen von §58 Abs.1 Satz 2 BNatSchG.  
[http://www.bsh.de/de/Meeresnutzung/Wirtschaft/Windparks/Windparks/Grundlagen/Leitsaetze\\_Eingriffsregelung.pdf](http://www.bsh.de/de/Meeresnutzung/Wirtschaft/Windparks/Windparks/Grundlagen/Leitsaetze_Eingriffsregelung.pdf) (zuletzt besucht am 08.12.2011)
- CAMPHUYSEN, K., PEET, G. & Maas, F.-J. (2006): Whales and Dolphins of the North Sea. – 's-Graveland/ Niederlande (Fontaine Uitgevers): 159 S.

- CARSTENSEN, J., HENRIKSEN, O. D. & TEILMANN, J. (2006): Impacts of offshore windfarm construction on harbour porpoises, acoustic monitoring of echolocation activity using porpoise detectors (T-PODs). – Marine Ecology Progress Series 321: 295-308.
- EVANS, P. G. H. (2008): Offshore wind farms and marine mammals: Impacts & methodologies for assessing impacts. – ECS Special Publication Series No. 49: 68 S.
- EVANS, P. G. H. & TEILMANN, J. (2009): Report of ASCOBANS/HELCOM small cetacean population structure workshop. – Bonn/ Deutschland (ASCOBANS): 141 S. [http://www.ascobans.org/sites/default/files/document/MOP6\\_5-08\\_PopulationStructureWorkshopReport\\_1.pdf](http://www.ascobans.org/sites/default/files/document/MOP6_5-08_PopulationStructureWorkshopReport_1.pdf) (zuletzt besucht am 08.12.2011).
- GILLES, A., SCHEIDAT, M. & SIEBERT, U. (2009): Seasonal distribution of harbour porpoises and possible interference of offshore wind farms in the German North Sea. – Marine Ecology Progress Series 383: 295-307.
- HAMMOND, P. S., BEARZI, G., BJØRGE, A., FORNEY, K., KARCZMARSKI, L., KASUYA, T., PERRIN, W. F., SCOTT, M. D., WANG, J. Y., WELLS, R. S. & WILSON, B. (2008): *Phocoena phocoena* (Baltic Sea subpopulation). – In: IUCN 2010. IUCN Red List of Threatened Species. Version 2010.4. – [www.iucnredlist.org](http://www.iucnredlist.org) (zuletzt besucht am 13.02.2011).
- JEFFERSON, T. A., Webber, M. A. & PITMAN, R. L. (2008): Marine Mammals of the World – A Comprehensive Guide to their Identification. – Amsterdam/ Niederlande (Academic Press): 573 S.
- JEPSON, P. D., BENNETT, P. M., DEAVILLE, R., ALLCHIN, C. R., BAKER, J. R. & LAW, R. J. (2005): Relationships between PCBs and health status of UK-stranded harbour porpoises (*Phocoena phocoena*). – Environmental Toxicology and Chemistry 24: 238-248.
- KANNAN, K., BLANKENSHIP, A. L., JONES, P. D. & GIESY, J. P. (2000): Toxicity reference values for the toxic effects of polychlorinated biphenyls to aquatic mammals. – Hum. Ecol. Risk Assess. 6: 181-201.
- KLEIVANE, L., SKAARE, J. U., BJØRGE, A., DE RUITER, E. & REIJNDERS, P. J. H. (1995): Organochlorine pesticide residue and PCBs in harbour porpoise (*Phocoena phocoena*) incidentally caught in Scandinavian waters. – Environmental Pollution 89: 137-146.
- KOCK, K.-H. & BENKE, H. (1996): On the by-catch of harbour porpoise (*Phocoena phocoena*) in German fisheries in the Baltic Sea and the North Sea. – Archive of Fishery and Marine Research 44: 95-114.
- KOSCHINSKI, S. & STREMPPEL, R. (2010): Strategien zur Vermeidung von Beifang von Seevögeln und Meeressäugetieren in der Ostseefischerei. – Neumünster (Nabu, GRD & GSM): 75 S.

- LAW, R. J., BERSUDER, P., MEAD, L. K. & JEPSON, P. D. (2008): PFOS and PFOA in the livers of harbour porpoises (*Phocoena phocoena*) stranded or bycaught around the UK. – Marine Pollution Bulletin 56: 770-797.
- LUCKE, K., LEPPER, P. A., HOEVE, B., EVERAARTS, E., V. ELK, V. & SIEBERT, U. (2007): Perception of low-frequency acoustic signals by a harbour porpoise (*Phocoena phocoena*) in the presence of simulated offshore wind turbine noise. – Aquatic Mammals 33: 55-68.
- MACLEOD, C. D., SANTOS, B. M., REID, R. J., SCOTT, B. E. & PIERCE, G. J. (2007): Linking sandeel consumption and the likelihood of starvation in the Scottish North Sea: could climate change mean more starving porpoises? – Biology Letters – doi: 10.1098/rsbl.2006.0588
- MADSEN, P. T., WAHLBERG, M., TOUGAARD, J., LUCKE, K. & TYACK, P. (2006): Wind turbine underwater noise and marine mammals: implications of current knowledge and data needs. – Marine Ecology Progress Series 309: 279-295.
- MEINIG, H., BOYE, P. & HUTTERER, R. (2009): Rote Liste und Gesamtartenliste der Säugetiere (Mammalia) Deutschlands. – Naturschutz und Biologische Vielfalt (Bonn/Deutschland) 70 (1): 115-153.
- PALKA, D. L., ROSSMAN, M. C., VANATTEN, A. S. & ORPHANIDES, C. D. (2008): Effect of pingers on harbour porpoises (*Phocoena phocoena*) bycatch in the US Northeast gillnet fishery. – Journal of Cetacean Research and Management 10 (3): 217-226.
- PALMÉ, A., LAIKRE, L. & RYMAN, N. (2004): Population genetics of harbour porpoise in Swedish waters – a literature review. – Naturvårdsverket report 5419: 53 S.
- READ, A. J. (1999): Harbour Porpoise *Phocoena phocoena* (Linnaeus, 1758). – In: RIDGWAY, S. H. & HARRISON, Sir R. (Hrsg.): Handbook of Marine Mammals, Vol. 6. – San Diego/ USA (Academic Press): 323-355.
- READ, A. J. & HOHN, A. A. (1995): Life in the fast lane: the life history of harbor porpoises from the Gulf of Maine. – Marine Mammal Science 11: 423-440.
- SANTOS, M. B. & PIERCE, G. J. (2003): The diet of harbour porpoise (*Phocoena phocoena*) in the northeast Atlantic. – Oceanography and Marine Biology: an Annual Review 41: 355-390.
- SCHEIDAT, M., KOCK, K.-H. & SIEBERT, U. (2004): Summer distribution of harbour porpoise (*Phocoena phocoena*) in the German North Sea and Baltic Sea. – Journal of Cetacean Research and Management 6 (3): 251-257.
- SCHEIDAT, M., GILLES, A. & SIEBERT, U. (2006): Evaluating the distribution and density of harbour porpoises (*Phocoena phocoena*) in selected areas in German waters. – In: VON NORDHEIM, H., BOEDEKER, D. & KRAUSE, J. C. (Hrsg.):

- Progress in Marine Conservation in Europe. – Berlin & Heidelberg/ Deutschland (Springer-Verlag): 189-208.
- SCHEIDAT, M., GILLES, A., KOCK, K.-H. & SIEBERT, U. (2008): Harbour porpoise *Phocoena phocoena* abundance in the southwestern Baltic Sea. – Endangered Species Research 5: 215-223.
- SKÓRA, K. E. & KUKLIK, I. (2003): Bycatch as a potential threat to harbour porpoises (*Phocoena phocoena*) in Polish Baltic waters. – NAMMCO Scientific Publications 5: 303-315.
- SOUTHALL, B. L., BOWLES, A. E., ELLISON, W. T., FINNERAN, J. J., GENTRY, R. L., GREENE, C. R., KASTAK, D., KETTEN, D. R., MILLER, J. H., NACHTIGALL, P. E., RICHARDSON, W. J., THOMAS, J. A. & TYACK, P. L. (2007): Marine mammal noise exposure criteria: Initial scientific recommendations. – Aquatic Mammals 33 (4): 411-521.
- STRAND, J., LARSEN, M. M. & LOCKYER, C. (2005): Accumulation of organotin compounds and mercury in harbour porpoises (*Phocoena phocoena*) from the Danish waters and West Greenland. – Science of the Total Environment 350: 59-71.
- SVEEGAARD, S. (2011): Spatial and temporal distribution of harbour porpoises in relation to their prey. – PhD thesis, National Environmental Research Institute. – Aarhus/ Dänemark (Aarhus University): 128 S.
- SVEEGAARD, S., TEILMANN, J., TOUGAARD, J., DIETZ, R., MOURITZEN, K. N., DESPORTES, G. & SIEBERT, U. (2011): High-density areas for harbor porpoises (*Phocoena phocoena*) identified by satellite tracking. – Marine Mammal Science 27 (1): 230-246.
- TOUGAARD, J., CARSTENSEN, J., TEILMANN, J., SKOV, H. & RASMUSSEN, P. (2009): Pile driving zone of responsiveness extends beyond 20 km for harbor porpoises (*Phocoena phocoena* (L.)). – Journal of the Acoustical Society of America 126 (1): 11-14.
- VERFUß, U. K., HONNEF, C. G. & BENKE, H. (2006): Seasonal and geographical variation of harbour porpoise (*Phocoena phocoena*) habitat use in the German Baltic Sea monitored by passive acoustic methods (PODs). – In: VON NORDHEIM, H., BOEDEKER, D. & KRAUSE, J. C. (Hrsg.): Progress in Marine Conservation in Europe. – Berlin & Heidelberg/ Deutschland (Springer-Verlag): 209-224.
- VERFUß, U. K., HONNEF, C. G., MEDING, A., DÄHNE, M., MUNDRY, R. & BENKE, H. (2007): Geographical and seasonal variation of harbour porpoise (*Phocoena phocoena*) in the German Baltic Sea revealed by passive acoustic monitoring. – Journal of the Biological Association of the United Kingdom 87: 165-176.
- VIAUD-MARTÍNEZ, K. A., MARTÍNEZ VERGARA, M., GOL'DIN, P. E., RIDOUX, V., ÖZTÜRK, A. A., ÖZTÜRK, B., ROSEL, P. E., FRANTZIS, A., KOMNENOU, A. & BOHONAK, A. J. (2007): Morphological and genetic differentiation of the Black Sea harbour

porpoise *Phocoena phocoena*. – Marine Ecology Progress Series 281: 294-338.

VINTHER, M. & LARSEN, M. (2004): Updated estimates of harbour porpoise (*Phocoena phocoena*) bycatch in the Danish North Sea bottom-set gillnet fishery. – Journal of Cetacean Research and Management 6: 19-24.

WIEMANN, A.; ANDERSEN, L. W., BERGGREN, P., SIEBERT, U., BENKE, H. & 9 WEITERE AUTOREN (2010): Mitochondrial control region and microsatellite analyses on harbour porpoise (*Phocoena phocoena*) unravel population differentiation in the Baltic Sea and adjacent waters. – Conservation Genetics 11: 195-211.

### Weiterführende Literatur und Links

BRÄGER, S. (2011): Die Ostsee-Schweinswale im Spannungsfeld des Artenschutzes. – Meer und Museum 23: 57-66.

HAELTERS, J. & CAMPHUYSEN, K. (2009): The harbour porpoise in the southern North Sea. – Ostend/ Belgium (IFAW): 56 S.

KOSCHINSKI, S. (2002): Current knowledge on harbour porpoises (*Phocoena phocoena*) in the Baltic Sea. – Ophelia 55 (3): 167-197.

NLWKN (Hrsg.) (2011): Vollzugshinweise zum Schutz von Säugetierarten in Niedersachsen. – Säugetierarten des Anhangs II der FFH-Richtlinie – Schweinswal (*Phocoena phocoena*). – Niedersächsische Strategie zum Arten- und Biotopschutz, Hannover, 13 S.

[http://www.nlwkn.niedersachsen.de/naturschutz/staatliche\\_vogelschutzwaerte/vollzugshinweise\\_arten\\_und\\_lebensraumtypen/vollzugshinweise-fuer-arten-und-lebensraumtypen-46103.html](http://www.nlwkn.niedersachsen.de/naturschutz/staatliche_vogelschutzwaerte/vollzugshinweise_arten_und_lebensraumtypen/vollzugshinweise-fuer-arten-und-lebensraumtypen-46103.html) (zuletzt besucht am 05.08.2013)

REID, J. B., EVANS, P. G. H. & NORTHRIDGE, S. P. (2003): Atlas of cetacean distribution in north-west European waters. – Peterborough/UK (Joint Nature Conservation Committee): 52-53.

SCHULZE, G. (1987): Die Schweinswale. – Neue Brehm Bücherei – Wittenberg Lutherstadt/ Deutschland (A. Ziemsen Verlag): 167 S.

SCHULZE, G. (1991): Wale an der Küste von Mecklenburg-Vorpommern. – Stralsund (Deutsches Meeresmuseum). – Meer und Museum 7: 22-52.

VAN DER MEIJ, S. E. T. & CAMPHUYSEN, C. J. (2006): The distribution and diversity of whales and dolphins (Cetacea) in the southern North Sea: 1970-2005. – Lutra 49 (1): 3-28.